

# **AQUATIC PLANT MANAGEMENT PLAN**

**FISH, CRYSTAL and INDIAN LAKES  
Lower Wisconsin River Basin**

**DANE COUNTY, WISCONSIN**



SCUBA cut channels through Eurasian watermilfoil in Fish Lake, 1990

January 2007

**OFFICE OF LAKES AND WATERSHEDS  
DANE COUNTY LAND AND WATER RESOURCES DEPARTMENT  
1 FEN OAK COURT, RM 234  
MADISON, WI 53718**

**Prepared by**

**Underwater Habitat Investigations, LLC**  
(Dave Marshall)

**and**

**Dane County Land & Water Resources  
and Public Works Staff**  
(Kevin Connors, Deb Flanders, Teresa Haynes, Sue Jones,  
Jim Leverance, Darren Marsh, Michelle Richardson, Joe Yaeger)

**Edited by  
Sue Jones**

**Cover photo by Dick Lillie**

---

**Preparation of this plan was funded in part by a Wisconsin  
Department of Natural Resources Lake Planning Grant**

**This plan was approved by the Wisconsin Department of Natural Resources on  
March 17, 2007 and by the Dane County Lakes and Watershed Commission  
on April 12, 2007**

## Summary

Aquatic plant surveys were performed during the summer of 2006 on Fish, Crystal and Indian lakes. The three lakes share common features including location within the Lower Wisconsin River Drainage Basin, status as seepage lakes, and occurrence of rising water levels in recent decades. Six hundred-ninety-three sites were sampled in the three lakes. Results of the point intercept surveys indicate that coontail and Eurasian watermilfoil dominate in both Fish and Indian lakes. Dense concentrations of these weedy plants coincide with low numbers of native species (low native species richness). In Fish Lake, a higher diversity of native plants went undetected since the sampling points did not capture the narrow band of nearshore plants. In Crystal Lake, Eurasian watermilfoil is a minor component in an aquatic plant community already limited by heavy blue-green algae blooms and poor water clarity. Plant density measures are lower in Crystal Lake compared to the other two lakes. Survey results indicate that Fish and Indian lakes warrant management of coontail and Eurasian watermilfoil while modest rooted plant densities in Crystal Lake are not posing water use problems. Higher water levels in both Indian and Crystal lakes play an important role in restoring sportfish populations due to fewer winterkills and improved habitat.

In addition to monitoring aquatic plant communities and assessing management strategies, public participation was planned and executed at all stages of plan development. Prior to conducting the aquatic plant surveys, two public meetings were advertised and conducted to educate the public on ecological and habitat values of aquatic plants and management issues. A final public meeting was held in December 2006 to share results of the aquatic plant surveys and draft aquatic plant management plan for public review and input. Public comment on the draft plan was solicited on the Dane County Office of Lakes and Watersheds/Lakes and Watershed Commission website.

Also, an Aquatic Plant Management Committee created by the Dane County Board met for much of the time coincident with plan field work and preparation. Members of that Committee were invited to provide comments on the draft plan. Recommendations from the Committee's final report (October 20, 2006) are included in Appendix A. This plan

will be amended as necessary to implement recommendations that are approved by the County Executive and County Board and implemented by county staff.

Ultimately, the plan recommendations do not represent a significant change from the relatively long history of mechanical harvesting operations on the lake. However, the plan identified important steps that will enhance the county's aquatic plant harvesting program and protect important habitat features in the lake.

### **Fish Lake Recommendations**

1. Conduct a more complete inventory of native aquatic plants near shore to better assess their status and condition following the first year of pumping and lake drawdown.
2. Conduct mechanical harvesting at important access points to improve navigation and fish habitat.
3. Mechanical and chemical removal of aquatic plants within Department of Natural Resources designated Sensitive Areas should be prohibited to protect critical habitat for fish spawning and early life stages.
4. Consider longer term efforts to sustain boating lanes and improved fish habitat using methods such as deep cutting - harvesting. Methods could include modified large scale harvesting or manual cutting involving SCUBA.
5. Protect important habitat features including floating-leaf plant beds and coarse woody habitat. Residents should be discouraged from manually removing high value species such as watershield, floating-leaf pondweed and water lilies.
6. Recommend Sensitive Areas Designations to WDNR based on criteria established in Wisconsin Administrative Code NR 107 and other important ecological features. Sensitive Areas would encompass plant beds with high value native species including watershield, floating-leaf pondweed and water lilies. Use of herbicides and large-scale mechanical harvesting is prohibited in these areas.
7. Encourage local land use planning and management to reduce nutrient runoff into the lake. (Watershed runoff contributed to littoral zone sediments rich in nutrients, a factor contributing to high Eurasian watermilfoil growth in the lake. Potential sources of polluted runoff should be re-evaluated given reductions linked to surrounding park land acquisitions.)

8. Sample nearshore fish populations, including blackchin shiner, blacknose shiner and banded killifish. These species may be affected by rapid habitat changes including rising water levels.
9. Update the comprehensive lake management plan at some point in the near future.
10. Dane County mechanical harvesting crews should continue to take steps to prevent the spread of exotic invaders across Dane County lakes. These steps include removing any visible plants, mud, debris, water, fish or animals from the machinery and thoroughly washing the equipment. The fact sheet in Appendix B is included in the harvesting crews' operations manual.

### **Crystal Lake Recommendations**

1. Mechanical harvesting should be conducted during periods when Eurasian watermilfoil densities are high to improve boating access.
2. Modest levels of native macrophytes (aquatic plants) provide important fish habitat and should not be eradicated. These conditions may change and Eurasian watermilfoil could expand under different water level conditions, warranting management.
3. Recommend Sensitive Area designations to WDNR including bays supporting white water lily beds.
4. Protect coarse woody habitat around the lake for fish and herptile populations.
5. Encourage local land use planning and management to reduce nutrient loading into the lake. (Reducing blue-green algae blooms could ultimately improve native plant growth in the lake.)
6. Consider coordinating the preparation of a comprehensive lake management plan with Columbia County.
7. Dane County mechanical harvesting crews should continue to take steps to prevent the spread of exotic invaders across Dane County lakes. These steps include removing any visible plants, mud, debris, water, fish or animals from the machinery and thoroughly washing the equipment. The fact sheet in Appendix B is included in the harvesting crews' operations manual.

## **Indian Lake Recommendations**

1. Continue harvesting channels when coontail or Eurasian watermilfoil impede navigation within the lake.
2. Avoid management activities that would shift productivity from submersed macrophytes to blue-green algae blooms. For example, massive primary productivity shift from rooted plants to blue-green algae blooms would likely result from whole-lake herbicide(s) treatment.
3. Consider planting floating leaf plants such as white water lily, yellow water lily or American lotus to improve fish habitat and aesthetics.
4. Sensitive Areas designation is not a high priority for Indian Lake at this time given the lake is entirely encompassed by park land, proposed management activities do not alter nearshore habitat, and high value aquatic plant species are very scarce in the lake.
5. Dane County mechanical harvesting crews should continue to take steps to prevent the spread of exotic invaders across Dane County lakes. These steps include removing any visible plants, mud, debris, water, fish or animals from the machinery and thoroughly washing the equipment. The fact sheet in Appendix B is included in the harvesting crews' operations manual.

## **Introduction**

As required in Wisconsin Administrative Code NR 109.04(d), the purpose of this plan is to guide mechanical harvesting activities and the effective management of aquatic plants in Fish, Crystal and Indian lakes. Dane County periodically operates mechanical harvesters in all three lakes, managing densely growing exotic Eurasian watermilfoil (*Myriophyllum spicatum*), exotic curly-leaf pondweed (*Potamogeton crispus*) and native coontail beds (*Ceratophyllum demersum*). Dense stands of these “weedy” plants undermine the ecological balance and recreational uses in these lakes. Harvesting efforts are designed to enhance both of these important lake management functions.

Significant ecological and hydrological changes occurred in all three lakes over the last several decades, altering aquatic plant distributions and species compositions. Long term rising water levels resulted in emergent, floating-leaf and submersed plant re-distributions while effects of polluted runoff benefited Eurasian watermilfoil and curly-leaf pondweed.

In recent years submersed aquatic plant growths expanded in the shallow hyper-eutrophic Indian and Crystal lakes, although native species richness remains low. In contrast, native plant beds declined significantly in Fish Lake, a deep water seepage lake with a maximum depth of 62 feet. Native plant declines in Fish Lake coincided with eutrophication, Eurasian watermilfoil invasion and water level changes.

The primary goals of this plan are to establish long term realistic objectives for managing nuisance exotic plant species while protecting valuable native species and their important habitat functions. While the goal is not to create a comprehensive lake management plan, aquatic plant communities are linked to other aspects of lake and watershed management, and these links are identified. A comprehensive lake management plan already exists for Fish Lake and this report updates and expands the aquatic plant management element of that plan.

While the plan is designed to develop a variety of strategies for managing aquatic plants, chemical treatments are not performed or sponsored by either Dane County or Wisconsin Department of Natural Resources (WDNR). WDNR regulates chemical herbicide treatments in public waters and permits are required under Administrative Code NR 107.


### **Fish Lake Goals**

Recognizing that Eurasian watermilfoil has dominated the littoral zone for over two decades, the goals for managing Fish Lake aquatic plants are to: (1) improve recreational access in the lake, (2) improve bluegill (*Lepomis macrochirus*) – largemouth bass (*Micropterus salmoides*) interactions within monotypic stands of Eurasian watermilfoil, (3) identify and protect Sensitive Areas defined under Wisconsin Administration NR 107.05(3-i) and (4) reverse the documented declines of high value species [NR 107.08(4)] in Fish Lake including large-leaf pondweed (*Potamogeton amplifolius*) clasping-leaf pondweed (*P. richardsonii*), sago pondweed (*Struckenia pectinatus*) and watershield (*Brasenia schreberi*). Other important native plants that declined in Fish Lake and also require protection include floating-leaf pondweed (*P. natans*), variable pondweed (*P. gramineus*), flat-stem pondweed (*P. zosteriformis*), smartweed (*Polygonum amphibian*), water marigold (*Bidens beckii*), yellow water lily (*Nuphar variegatum*), white water lily (*Nymphaea tuberosa*), and American lotus (*Nelumbo lutea*).

## Background Information

Fish Lake (252 acres) is a moderately eutrophic lake located in the Town of Roxbury (Figure 1). The lake is relatively undeveloped with significant parklands adjoining the east and west shorelines. Recreational uses include swimming, fishing and boating. There is a town ordinance prohibiting gasoline motors on the lake.

Figure 1: Fish, Crystal and Indian lakes location within Dane County



The Fish Lake watershed is approximately 1680 acres including lake surface areas. The primary land use is agriculture. Top soils are fine silty loam and are nutrient rich from manure and fertilizer applications. Most of the watershed is rolling farmland with steep wooded hills. Just northwest of Fish Lake is Mud Lake (74 acres), historically a northwest bay of Fish Lake that was partly disconnected when Fish Lake road was constructed. The bay is currently connected via a culvert.

Major changes occurred within Fish Lake over the last several decades including declining water quality and reduced native aquatic plant beds. Detailed information on Fish Lake can be found in a comprehensive lake management plan (Marshall et al. 1996) and in numerous research articles focusing on ecology of macrophytes and fish. The comprehensive lake management plan was based on a U.S. Environmental Protection Agency (USEPA) Clean Lakes Phase I Diagnostic and Feasibility Study and incorporated significant findings of the cooperative research effort known as the “Integrated Management of Macrophytes and Fish.”

Historically, Fish Lake was classified mesotrophic based on chlorophyll-a, phosphorus, and Secchi (Dane County Regional Planning Commission – DCRPC 1979). During the 1970’s, the lake had the best water quality in the county, however other indicators suggested gradual water quality decline. Hypolimnetic dissolved oxygen levels declined since the late 1950’s while poor survival of stocked rainbow trout (*Oncorhynchus mykiss*) ended any efforts to manage a two story fishery by 1969 (DCRPC 1979). Cisco (*Coregonus artedii*) populations are native to the lake, and like trout also require deep cool water habitat with sufficient dissolved oxygen. Over the past several decades, periodic cisco kills were documented and coincided with low dissolved oxygen levels in the upper hypolimnion and thermocline.

Fish Lake historically supported diverse floating-leaf and submersed aquatic plant beds (Table 1) but significant declines in abundance have occurred. Native plant declines coincided with three long-term changes in the lake: eutrophication, Eurasian watermilfoil (EWM) invasion and rising water levels.

As of the 1996 comprehensive lake management plan, approximately 60% of the Fish Lake watershed was agricultural and primarily in the form of cropland. Even though the watershed to lake ratio is relatively low at 4.4:1, high phosphorus loading was documented during the 1990’s. The annual estimated phosphorus loading to the lake was 1690 lbs/year based on lake modeling conducted as part of the comprehensive plan. Winter manure spreading and feedlots were identified as principal watershed sources of phosphorus and nitrogen at that time. More recently, the predicted phosphorus loading to the lake has declined and reflects a feedlot closure near Mud Lake and expanded parkland around both Fish and Mud lakes.

Within the last few decades, rising Trophic State Index (TSI) values indicated that Fish Lake shifted from mesotrophic to moderate eutrophic condition. The long-term water quality decline in the lake was linked to watershed nutrient sources (Marshall et al. 1996). Evidence of declining water quality included reduced Secchi measurements, higher chlorophyll and higher hypolimnetic phosphorus and ammonia levels. In addition to increasing (TSI) values, littoral zone sediments also reflected nutrient enrichment. Shallow water sediment core sampling revealed very high levels of both phosphorus (1142 mg/kg) and ammonia (128 mg/l). Sediment testing indicated that polluted runoff was deposited within littoral areas of the lake, particularly along the west shorelines adjacent to most of the agricultural runoff. Sediment fertility was linked with EWM growth, and phosphorus transport from the littoral zone likely occurred as well (Smith and Barko 1990). Deep water sediment core sampling was also conducted and revealed significant water quality decline in recent years. Analyzing sediment cores is a way of determining a history of nutrient input into a lake. Upper portions of sediments reflect recent deposition. Deeper sediment layers correspond with earlier deposition.

While detailed lake and watershed monitoring studies were initiated in 1988 to address the declining water quality, lake users were generally more aware of the “dense weed beds” in the lake. Eurasian watermilfoil was first identified in 1967 and rapidly expanded throughout the 1980s. By 1991 dense growths of EWM covered 99 acres of the lake bottom area (Lillie 1996). During the EWM expansion period, numerous native species declined substantially as EWM established monotypic stands beyond one meter depth - a typical pattern of EWM invasions (Madsen et al. 1991). With the exception of coontail, the remaining native macrophytes occupied near-shore areas (Lillie 1996). The near-shore native plant beds can be more vulnerable to shoreline development and rapid water level decline.

In 1994, EWM declined by approximately 40% across the lake. The decline coincided with weevil damage (Lillie 2000, Creed 1998). Native weevils can reduce the viability of EWM by boring into the stems (Mazzei et al. 1999). Boring into the stem results in loss of plant buoyancy and the plant sinks to the bottom. This either kills the plant directly or severely weakens the plant due to reduced photosynthesis. Coinciding with reduced aquatic plant density that year, Secchi depths declined and chlorophyll-a concentrations increased. Higher chlorophyll levels may have reflected nutrient release from decaying

EWM, reduced alleopathy or both. These conditions were temporary since EWM rebounded in 1996. The temporary EWM decline did not expand the distribution or abundance of native plants and may indicate that the high sediment nutrient levels favor EWM over environmentally sensitive native aquatic plants. The EWM decline and resurgence suggest that a lake-wide chemical eradication may not expand native plants and could result in severe blue-green algae blooms.

In addition to eutrophication and EWM expansion in Fish Lake, long-term rising water levels (Krohelski et al. 2002) was a third factor contributing to redistribution of native plants. As the water level rose, emergent and floating-leaf plants moved to newly submersed shoreline while EWM also migrated toward shore. The result was a gradual shift of all plants (emergent, floating-leaf and submersed) toward the perimeter of the lake. In 2006, the management district began pumping water from the lake to reduce water levels and damage to several properties along the west shore. Many of the relatively scarce native species became desiccated as water levels dropped.

The EWM invasion has altered the habitat chemically in Fish Lake (Unmuth et al. 2000). Very low dissolved oxygen levels were found near the bottom of the beds. The effects of dense plant beds on predator-prey interactions were reported as well (Engel 1987, Savino et al. 1992). Local efforts to develop new methods for improving habitat within dense EWM beds began in 1989 (Marshall 1990). SCUBA divers used manual cutting tools in Fish Lake to cut deeper growths of EWM at the sediment surface. The deep cutting technique held promise since the channels created by the SCUBA divers persisted for four years. Aerial photographs of the lake during this period clearly show where the channels were cut (cover photo). Modest growths of curly-leaf pondweed and coontail replaced EWM within the channels. Deep cutting to stress deeper EWM stands was ultimately tested by teams of researchers seeking management tools for improving EWM habitat and predator-prey interactions (Unmuth et al. 1999, Unmuth et al 1998, Olson et al. 1998, Trebitz et al 1997). Dane County Public Works staff modified one of the county harvesters in order to conduct a series of deep cutting experiments in Fish Lake and other lakes around the state. While the mechanical channels did not persist as long as the manual cut channels, the results demonstrated increased growth rates for particular year classes of both bluegill and largemouth bass populations. “Cruising lanes” became

available to largemouth bass. Predation on stunted bluegills occurred, followed with increased growth rates of specific year classes for both species.

Bluegill and largemouth bass comprise the dominant fisheries in the lake. Other species found in Fish Lake are listed in Table 2. Environmentally sensitive nongame species have been identified in Fish Lake, including banded killifish (*Fundulus diaphanous*), blackchin shiner (*Notropis heterodon*), blacknose shiner (*Notropis heterolepis*), and Iowa darter (*Etheostoma exile*). These species can typically be found in dense aquatic plant communities near shore (Becker 1983). The banded killifish is classified as State Special Concern and the other three species are classified as environmentally sensitive to degraded habitat (Lyons 1992). Abundant overhanging trees ring the lake and create another important habitat feature for fish populations and herptiles. In 2002, WDNR and Dane County Parks cooperated in a habitat improvement project along the Lussier Park shore. Large dead trees were pushed into the water and American lotus seed and nursery seedlings from Mud Lake were planted as well. The goal was to improve habitat for game fishes and intolerant nongame species that can be vulnerable to near-shore habitat loss. The current status of nongame fishes in the lake is unknown.

Table 1: Fish Lake floating-leaf and submersed macrophytes (Lillie 1996)

Scientific Name	Common Name
<i>Bidens beckii</i>	Water marigold
<i>Brasenia schreberi</i>	Water Shield*
<i>Ceratophyllum demersum</i>	Coontail*
<i>Chara</i>	Stonewort or Muskgrass
<i>Eleocharis acicularis</i>	Spike Rush
<i>Elodea canadensis</i>	Elodea
<i>Lemna minor</i>	Lesser Duckweed
<i>Myriophyllum sibiricum</i>	Northern Watermilfoil
<i>Myriophyllum spicatum</i>	Eurasian Watermilfoil (EWM – exotic)*
<i>Najas flexilis</i>	Bushy Pondweed
<i>Nuphar variegata</i>	Yellow Water Lily
<i>Nymphaea odorata</i>	White Water Lily*
<i>Polygonum natans</i>	Smartweed*
<i>Potamogetan amplifolius</i>	Large-leaf Pondweed
<i>Potamogetan crispus</i>	Curly-leaf Pondweed (CLP-exotic)*
<i>Potamogetan gramineus</i>	Variable-leaf Pondweed
<i>Potamogetan natans</i>	Floating-leaf Pondweed*
<i>Potamogetan richardsonii</i>	Clasping-leaf Pondweed*

<i>Potamogetan zosteriformes</i>	Flat-stem Pondweed
<i>Struckenia pectinatus</i>	Sago Pondweed*
<i>Utricularia vulgaris</i>	Bladderwort

\* Herbarium specimens collected in 2006.


Table 2: Fish taxa sampled from Fish Lake in 1991

Scientific Name	Common Name
<i>Esox lucius</i>	Northern pike
<i>Cyprinus carpio</i>	Common carp
<i>Notemigonus crysoleucas</i>	Golden shiner
<i>Notropis heterodon</i>	Blackchin shiner
<i>Notropis heterolepis</i>	Blacknose shiner
<i>Catostomus commersoni</i>	White sucker
<i>Ameiurus natalis</i>	Yellow bullhead
<i>Fundulus diaphanus</i>	Banded killifish
<i>Lepomis cyanellus</i>	Green sunfish
<i>Lepomis gibbosus</i>	Pumpkinseed sunfish
<i>Lepomis macrochirus</i>	Bluegill sunfish
<i>L. gibbosus</i> x <i>L. cyanellus</i>	Hybrid sunfish
<i>L. gibbosus</i> x <i>L. macrochirus</i>	Hybrid sunfish
<i>Micropterus salmoides</i>	Smallmouth bass
<i>Pomoxis nigromaculatus</i>	Black crappie
<i>Etheostoma exile</i>	Iowa darter
<i>Perca flavescens</i>	Yellow perch
<i>Stizostedion vitreum</i>	Walleye

### Recent Aquatic Plant Management and Harvesting Records

The primary focus of mechanical harvesting in Fish Lake is to improve boating access from the Fish Lake Park launch and along developments around the lake. Harvesting records from 1999 through 2004 indicate that plant tonnages removed from the lake ranged from 13 to 104 tons of plants ( 40 tons – 1999, 104 tons – 2000, 13 tons – 2001, 18 tons - 2004). Figure 2 documents Dane County harvesting records since 1989.

Figure 2: Fish Lake mechanical harvesting summary



## 2006 Aquatic Plant Survey Update

### Methods

The sampling protocol was developed by Jen Hauxwell, a research scientist with Wisconsin Department of Natural Resources, Bureau of Integrated Science Services. The point intercept method was used where a large number of sampling sites are distributed equidistantly across a lake. GPS units were used to locate the sites and double-headed rakes were used to collect macrophytes. The rakes were constructed in two forms. The pole rake was used for sampling macrophytes up to 15 ft (4.6 m) and rope rake can be used to sample deeper areas. Density ratings from 1-3 were determined by the amount of plant material in the two-headed rake. Plants that were observed near the boat but were not collected in the rake were also noted. Samples of each species found in a lake were collected, pressed and submitted as voucher specimens to the UW Madison Herbarium.

Statistical analysis included the following:

- Frequency of occurrence within vegetated sites (number of times a species was found divided by the total number of vegetated sites).
- Relative frequency of plant species collected (describes each species contributing a certain percentage of the whole aquatic plant community).

Detailed statistical results appear in Appendix C.

Wisconsin Department of Natural Resources provided the sampling grids and spreadsheet software for data entry and analysis. A more detailed sampling description can be found in *Baseline Monitoring of Aquatic Macrophytes* (Hauxwell 2006).

### **Results and Discussion**


The aquatic plant survey was conducted on August 1 and 4, 2006. Secchi measurements were 6 feet on both survey days, generating a TSI score of 52 (moderately eutrophic). A 394-point grid was established across the lake containing potential sampling sites (Figure 3). After eliminating sites within depths well beyond the littoral zone, a total of 226 points were sampled.

Aquatic plant sampling depths ranged from 5 feet to 36 feet with a mean depth of 14.4 feet. The grid system was created from an older map and did not capture shallow nearshore areas around the lake. The sampling depths in Fish Lake were generally greater than other Dane County lakes sampled for macrophytes in 2006 due to the lack of nearshore sites and presence of moss beyond the littoral zone. Mosses were found to grow within the depth range (17 – 31 feet) typically defined as littoriprofundal (Wetzel 1983). Implications for the range of sampling depths were twofold. First, the minimum depth of 5 feet resulted in under sampling native plants that were located close to shore. Second, sampling moss beyond the typical littoral zone resulted in lower relative and vegetated site frequencies for both EWM and coontail (Figure 4). The results were adjusted in Figure 5 by removing the moss data. Regardless of these sampling issues, EWM and coontail dominated most of the littoral zone and reflected low native species richness beyond 5 feet water depth. Only one other species was found during the point intercept sampling effort. Actual plant distributions across Fish Lake are located on the maps in Appendix D. A single watershield specimen was collected along the west shoreline. The average rake fullness rating was 2 for both EWM and coontail.

Figure 3: Point intercept sampling grid for Fish Lake




Figure 4: Fish Lake vegetated site occurrence and relative frequency results



The relative frequency describes the percentage that each aquatic plant species contributes to the whole aquatic plant community. Relative frequency is a commonly used metric since survey results can be compared with surveys that used different sampling techniques.


Figure 5: Fish Lake vegetated site occurrence and relative frequencies without moss



The relative frequency describes the percentage that each aquatic plant species contributes to the whole aquatic plant community. Relative frequency is a commonly used metric since survey results can be compared with surveys that used different sampling techniques.

The sampling grid provided by DNR (Figure 3) did not include the southwest bay but observations confirmed dominance of EWM and coontail in that area as well. Both species were also dominant in that area when the lake was sampled from 1991-94. Because the sampling grid did not include nearshore areas during the 2006 survey, native plants and herbarium specimens closer to shore were periodically collected and identified. While the additional specimen collections did not constitute a complete nearshore sampling effort, the presence of white water lily, smartweed, clasping-leaf pondweed, floating-leaf pondweed, and sago pondweed were identified at various locations around the lake. Sampling planned for summer 2007 will both expand the point intercept sampling sites within the southwest bay and elsewhere, and include a more thorough survey of nearshore plants. This plan will then be amended to incorporate those results.

Figure 6: Fish Lake native species richness – 2006 survey, 2006 observations, and 1991-94 surveys



American lotus was observed growing in abundance around adjoining Mud Lake. No evidence was found of American lotus in Fish Lake including the Lussier Park area where it was planted in 2002 as part of a habitat restoration effort. Several species including American lotus, white water lily, smartweed, clasping-leaf pondweed, and sago pondweed were found desiccated within the drawdown zone of both Fish and Mud Lakes. In addition to the nearshore native plant observations, a few curly-leaf pondweed (CLP) turions were observed floating, an indication that the plant was in the over-wintering stage during the survey.

Native species richness was greater during the more intensive macrophyte surveys from 1991-94, perhaps reflecting different sampling methodologies and more intensive nearshore sampling at that time. Figure 6 compares native species richness data for the 2006 point intercept sampling effort, 2006 combined sampling with observations, and 1991-94 surveys results. Additional nearshore sampling is required to determine if a decline in native species richness has occurred. Aside from these differences, the general pattern of EWM dominating throughout the deeper portion of the littoral zone did not change. Coontail was the second most abundant plant during all of the surveys.

Sensitive Areas (NR 107.05(3)(i)) were described in the 1996 comprehensive management plan. Areas supporting watershield, floating-leaf pondweed, white water lily, and yellow water lily included the west shore, southwest bay, northwest area, and east shoreline. Based on the 2006 aquatic plant survey, these areas continue to provide critical habitat for fisheries, including the environmentally sensitive nongame fishes and early life stages of sportfish in the lake. The floating-leaf plant communities should not be chemically treated or mechanically harvested. Small manual aquatic plant removal should be limited to Eurasian watermilfoil, curly-leaf pondweed and coontail. The proposed Sensitive Areas are identified in the Figure 7 map.

Appendix E notes fish and waterfowl values of desirable native plants in Fish, Crystal and Indian lakes.

Figure 7: Proposed Fish Lake sensitive areas



## **Aquatic Plant Management Alternatives**

While the primary emphasis of this plan is to protect important aquatic plant habitats and control nuisance EWM growths with mechanical harvesting equipment, additional management tools are available to individual property owners. Chemical treatments are regulated under Wisconsin Administrative Code NR 107. WDNR has generally discouraged chemical treatments in the past due to the vulnerability of nearshore native plant communities. Instead, riparian property owners have been encouraged to manually remove vegetation to enhance private access in front of their properties. Under NR 109.06 (a-1), a riparian owner is not required to obtain a permit from WDNR if the removal involves invasive species or removal of native species is limited to a single area with a maximum width of no more than 30 feet measured along the shoreline. In general, chemical treatments may not provide long-term EWM control in Fish Lake due to high littoral zone sediment nutrients.

Creating long-term access channels in Fish Lake could involve either deep cutting and harvesting or hydraulic dredging. Due to high costs, dredging may not be a practical option. This form of aquatic plant management would also require a Chapter 30 permit from WDNR. As prior research has demonstrated, deep cutting and harvesting can sustain relatively long-term access channels. This effort may involve modification of existing mechanical harvesters or labor intensive efforts such as manual cutters or participation of SCUBA divers.

Another alternative is the use of aquatic weevils. Weevils have been demonstrated to control EWM in laboratory and enclosure studies (Mazzei et al. 1999, Sheldon and Creed 1995). The EWM decline in 1994 coincided with evidence of weevil damage (Lillie 2000), however EWM rebounded a few years later and high densities continue in the presence of the insect. More detailed discussions on aquatic plant management alternatives can be found in Cooke et al. (2005) and Petty (2005).

Mechanical harvesting often removes juvenile panfish in the conveyor with the vegetation. However, WDNR biologists consider the inadvertent removal of small panfish to have no effect on population levels. In some cases, removal of juvenile panfish may actually improve size structure when population levels are too high.

## Specific Alternatives

- 1) **No treatment:** Rejecting all types of aquatic plant management does not appear realistic, given the extent of EWM coverage. Severe ecological unbalance already exists as well as significant loss of recreational uses over approximately 40% of the surface area.
- 2) **Biological control:** This method does not appear realistic at this time. Weevil (*Euhrychiopsis lecontei*) populations have already been well documented in Fish Lake, yet the density of EWM declined significantly only once over the past two decades. The findings are not clear if weevils were the cause for the temporary decline in 1994. High panfish populations can also reduce weevil densities below levels needed for effective EWM control.
- 3) **Chemical control:** Herbicide use should be restricted to small riparian areas using agents selective at controlling EWM. 2, 4-D is the likely agent given the partial selectivity for controlling EWM. However, several valuable native plants including water lilies can be damaged from 2, 4-D so WDNR permit applications should be carefully screened to avoid loss of already declining native plants. Whole-lake applications do not appear realistic given the past EWM decline and resurgence that may reflect sediment, nutrients and other factors.
- 4) **Manual - hand removal:** Manually removing plants around piers and swimming areas is a viable option. However, property owners should be educated about the importance of high value native species so that their efforts should focus on weedy exotics such as EWM. Sensitive Areas cannot be harvested. SCUBA divers can be used for larger areas such as creating long-term boat channels within the manual removal zone. All plants that are cut must be removed. In the case of SCUBA divers, barges or pontoon boats can be used to transport harvested plants from the lake.
- 5) **Mechanical harvesting:** Given the extent of EWM over 40% of the lake surface area, mechanical harvesting provides effective temporary access through the dense monotypic beds as well as habitat improvements. Deep cutting and harvesting offers potential for relatively long-term effectiveness but modifications to existing harvesters would be required.
- 6) **Physical control:** Most physical methods of controlling aquatic plant require permits from WDNR under Chapter 30 Wisconsin State Statutes. Hydraulic dredging can be an option for removing the nutrient-rich sediments within

designated navigation channels. This method has the greatest potential for long-term control but can be initially expensive. Whole lake dredging is unrealistic given the vast littoral areas affected by EWM. Sheets of dark fabrics anchored to the bottom covering the plants from any light create bottom or benthic barriers. A benthic barrier eliminates all plants, including non-target species, with the exception of free-floating plants and algae. These types of barriers can interfere with fish spawning and other pond wildlife. Over time, barriers may require maintenance due to siltation, ice damage, bubbling up and normal wear and tear. Fabrics applied in this way will require a permit. Fabrics are rarely used by property owners because of the labor of installation and maintenance. During local demonstrations, problems arose due to gas collection under the fabric. Drawdown is also rarely used in Wisconsin for aquatic plant management, however drawdown is currently underway in Fish Lake with the goal of water level control and not aquatic plant management. So far negative effects on valuable native plants have been observed.

### **Crystal Lake Goals**

The goals for managing Crystal Lake macrophytes are to: (1) control dense EWM and other weedy plant growths in areas where recreational uses are undermined, (2) identify and protect Sensitive Areas defined under Wisconsin Administration NR 107.05(3-i), particularly floating-leaf plant beds, and (3) sustain native aquatic plant beds for fish habitat and inhibit potentially toxic concentrations of blue-green algae blooms that frequently occur within this hypereutrophic lake.

### **Background Information**

Crystal Lake is a 525-acre shallow seepage lake located just 1,950 feet east of Fish Lake (Figure 1). Recreational uses include gasoline motorized boating, fishing, water skiing and swimming. In recent years, Crystal Lake has been a popular attraction for anglers due to the fast growing bluegill, crappie and largemouth bass populations. Additional recreational opportunities are located at a large commercial park located on the Columbia County side of the lake.

Unlike the relatively deep Fish Lake, Crystal Lake is shallow and it does not thermally stratify. Crystal Lake is classified as hypereutrophic due to high concentrations of blue-green algae. The WDNR lake database indicates that Secchi depth measurements range from 1.5 (TSI = 72) to 2.8 feet (TSI = 63). The surrounding watershed is very similar to the Fish Lake watershed with agriculture the dominant land use. Likely sources of phosphorus to Crystal Lake include feedlots, crop fields and internal loading as the lake mixes throughout the summer. During the 1980s, WDNR conducted animal waste management (NR 243) investigations on several shoreline feedlots located on the Columbia County side of the lake.

Crystal Lake is hydraulically connected to Fish Lake by groundwater, and rising water levels have coincided in both lakes for decades (Krohelski et al. 2002). Maximum water depths were only 6 feet in the 1940s and increased to 9 feet by 1960. Frequent winter fish kills had been documented from the 1940s through the 1960s (DCRPC 1979). Aeration and frequent stocking were necessary to create recreational fishing during that period. When fish kills occurred, bullheads were often the only survivors.

In recent years, the trend of increasing water levels continued and the maximum water depth increased to 14 feet. Consistent with the Fish Lake shoreline, trees became inundated in past years and dead trees now line the perimeter of Crystal Lake. The dead trees are an important habitat feature for fish and herptile populations. Coinciding with the rising water levels, sustainable largemouth bass and panfish populations in the lake indicate that winterkills have diminished. In spite of continued hypereutrophic conditions, greater water volume has apparently increased the total oxygen mass within the lake. Water level declines in the future, perhaps linked to pumping, may potentially reverse the trend of sustainable winter dissolved oxygen levels in the lake.

Dense growths of macrophytes were reported decades ago including common waterweed (*Elodea canadensis*), sago pondweed (*Struckenia pectinatus*), duckweed (*Lemna*), and white water lily (*Nymphaea odorata*) (DCRPC 1970). There are no historical quantitative records on blue-green algae blooms or how the blooms might have affected the maximum rooting depths and distribution of macrophytes in Crystal Lake.

In recent years, EWM became established in the lake and the formation of dense monotypic stands created recreational use problems. Management included private herbicide applications around the two commercial mobile home parks in Columbia County, while Dane County operated mechanical harvesters to provide boating access from the public boat ramp and elsewhere.

Fish populations have fluctuated over the years due to previous winterkills and also reflect restocking efforts. Bluegills (*Lepomis macrochirus*), largemouth bass (*Micropterus salmoides*), and black bullhead (*Ameiurus melas*) have been the most common species reported (DCRPC 1979). Other species reported from the lake include golden shiner (*Notemigonus crysoleucas*), fathead minnow (*Pimephales promelas*), pumpkinseed (*Lepomis gibbosus*), and orange spotted sunfish (*Lepomis humilis*). No environmentally intolerant fish species have been reported from the lake. Since about 1980, higher water levels coincided with sustainable populations of largemouth bass, bluegill, black crappie (*Pomoxis nigromaculatus*), and, to a lesser extent, yellow perch (*Perca flavescens*) (Larson – personal communication). In addition to high water levels, native macrophytes enhance fisheries habitat. Dissolved oxygen deficits, associated with low water level induced winterkills or heavy summer blue-green algae blooms, and outbreaks of *Columnaris* bacteria are factors that can have negative impacts on the fisheries.

## **2006 Aquatic Plant Survey Update**

**Methods** (Refer to page 14)

### **Results and Discussion**

The aquatic plant survey was conducted on July 28 and 29, 2006. Secchi measurements reflected very poor water clarity with depths ranging from 1.8 to 2.5 feet during the survey. The equivalent Secchi TSI scores were 70 and 63 respectively, clearly indicating eutrophic conditions. The original sampling grid contained 495 points (Figure 8) however the total number was reduced. There were more sampling points created than were needed to conduct the survey, given the typically low native species richness found in hypereutrophic lakes such as Crystal Lake. A total of 263 points were ultimately sampled but total sampling area was maintained since every other point was sampled along a given transect.

Consistent with the Fish Lake sampling grid, most of the sampling points were not very close to shore since the maps did not reflect the current lake volumes. For instance, in the Figure 8 grid sampling provided by DNR, a portion of the northeast bay was excluded. Historically, that area was partly a cattail marsh. Collection of specimens closer to shore was attempted, however nearshore native species richness did not appear to differ from deeper locations and the only additional plants observed were white water lily and cattail (*Typha*). Overall, the sampling grid did not significantly affect the plant native species richness score. Sampling planned for summer 2007 will capture areas excluded in 2006, and this plan will then be amended to incorporate those results.

Unlike Fish Lake, where EWM is clearly the dominant macrophyte, native species are dominant in Crystal Lake. Common waterweed (*Elodea canadensis*) is the most abundant plant followed by leafy pondweed (*Potamogeton foliosus*). Filamentous algal growths are also relatively abundant in the lake and are often attached to the aquatic plants. Both the vegetated sites frequency and relative frequency results reflect the dominance of common waterweed in Crystal Lake (Figure 9). Results also indicate that EWM was infrequently found in 2006 and it did not pose recreational or ecological problems. The average rake fullness was 1 for EWM and 2 for common waterweed, coontail and sago pondweed. Native species richness was actually higher in Fish Lake if the nearshore plants that were not part of the point intercept sampling effort are included. Appendix E notes fish and wildlife values of desirable native plant species in Fish, Crystal and Indian lakes. Distributions of the plants found in Crystal Lake are displayed in Appendix F.

Curly-leaf pondweed (*Potamogeton crispus*) is the other exotic plant found in the lake but at a very low frequency. Curly-leaf pondweed (CLP) typically becomes dormant when water temperatures increase and most late summer aquatic plant surveys either underestimate typical densities or miss the plant entirely. Windrows of CLP turions were observed floating across the lake during the Crystal Lake survey. High numbers of the over-wintering turions indicated potentially high densities of the plant earlier in the growing season. Frequency and distribution of CLP could be assessed more effectively if an early season survey was performed.

Sensitive Areas have not been designated on Crystal Lake. Based on the 2006 survey results, Sensitive Areas should include portions of the northeast bay and west shoreline (Figure 10) that support white water lilies. Chemical treatments and large-scale mechanical harvesting should be avoided in these areas.

Figure 8: Point intercept sampling grid for Crystal Lake





Figure 9: Crystal Lake vegetated site occurrence and relative frequency results



The relative frequency describes the percentage that each aquatic plant species contributes to the whole aquatic plant community. Relative frequency is a commonly used metric since survey results can be compared with surveys that used different sampling techniques.

Figure 10: Proposed Crystal Lake sensitive areas



## Recent Aquatic Plant Management

Dane County records indicate that mechanical harvesting has not been performed on the lake since 1999. The combination of relatively low densities of rooted plants and dense blue-green algae blooms have eliminated the need for harvesting in recent years. Chemical herbicide applications have been conducted within the vicinity of the mobile home parks on the northwest side of the lake.

## Aquatic Plant Management Alternatives

Results of the 2006 aquatic plant survey indicate that overall macrophyte densities are low in Crystal Lake. Macrophyte densities are notably low within the northwest bay area that received chemical treatment in 2006 and previous years. The overall low densities suggest that major macrophyte control efforts are not warranted in the lake. Rather, the macrophytes densities generally provide desirable habitat functions for fish. If a resurgence of EWM is detected in the lake, then a combination of selective chemical applications and mechanical harvesting beyond the nearshore areas may improve habitat and recreational uses. Given the hypereutrophic condition of the lake, continued blue-green algae blooms will likely continue in the lake. The long-term challenge is to reduce nuisance growths of exotic CLP and EWM while enhancing native populations that may compete with blue-green algae and provide habitat for fish.

Mechanical harvesting often removes juvenile panfish in the conveyor with the vegetation. However, WDNR biologists consider the inadvertent removal of small panfish to have no effect on population levels. In some cases, removal of juvenile panfish may actually improve size structure when population levels are too high.

### Specific Alternatives

- 1) **No treatment:** Rejecting all types of aquatic plant control may be reasonable given the modest densities of both native aquatic plants and EWM.
- 2) **Biological control:** This method does not appear realistic at this time. Weevil (*Euhrychiopsis lecontei*) populations had been well documented in Fish Lake, yet the density of EWM declined significantly only once over the past two decades.

The findings are not clear if weevils were indeed responsible for the temporary decline in 1994. EWM densities currently are not causing ecological or habitat problems in the lake.

- 3) **Chemical control:** Herbicide use should be restricted to small riparian areas and commercial campgrounds using agents selective at controlling EWM. 2,4-D is the likely agent given the partial selectivity for controlling EWM. However, several valuable native plants including water lilies can be damaged from 2,4-D so permit applications should be carefully screened to avoid loss of already declining native plants. Whole-lake applications do not appear feasible given the low density of EWM and frequent blue-green algae blooms.
- 4) **Manual - hand removal:** Manually removing plants around piers and swimming areas is a viable option. However, property owners should be educated about the importance of high value native species so that their efforts focus on weedy exotics such as EWM. All plants that are cut must be removed. Sensitive Areas cannot be harvested.
- 5) **Mechanical harvesting:** In 2006, mechanical harvesting was not needed due to the low densities of aquatic plants. However, this option should remain available if a resurgence of EWM undermines lake access.
- 6) **Physical control:** Most physical methods of controlling aquatic plant require permits from WDNR under Chapter 30 Wisconsin State Statutes. Hydraulic dredging is an unlikely option in Crystal Lake. The heavy blue-green algae blooms will likely replenish sediment nutrients within a short period following excavation. Watershed nutrient sources are likely an issue as well. Sheets of dark fabrics anchored to the bottom covering the plants from any light create bottom or benthic barriers. A benthic barrier eliminates all plants, including non-target species, with the exception of free-floating plants and algae. These types of barriers can interfere with fish spawning and other pond wildlife. Over time, barriers may require maintenance due to siltation, ice damage, bubbling up and normal wear and tear. Fabrics applied in this way will require a permit. Fabrics are rarely used by property owners because of the labor of installation and maintenance. In local demonstrations, problems arose due to gas collection under the fabric. Drawdown is currently underway in Fish Lake, and Crystal Lake water levels are predicted to decline due to the hydraulic connection. While the goal is not aquatic plant

management but rather flood control, valuable nearshore native plants can be negatively affected.

### **Indian Lake Goals**

The goals for managing Indian Lake macrophytes are to: (1) improve non-motorized boat access within dense coontail, Eurasian watermilfoil and curly-leaf pondweed beds, (2) sustain lake-wide aquatic plant beds in desirable densities to prevent blue-green algae blooms that had historically occurred, (3) manage aquatic plants to enhance the largemouth bass and bluegill fisheries, and (4) enhance native floating-leaf plant populations.

### **Background Information**

Indian Lake is a 27 ha (66 acres) shallow kettle lake that is maintained by groundwater and surface runoff (see Figure 1 for location). The entire lake is surrounded by the county park, and recreational uses include fishing, bird watching, canoeing and other types of boating that do not involve gas engines. The lake is primarily managed for largemouth bass and panfish. An aeration system is frequently used during late winter months to avoid anoxia and fish winterkill conditions.

The small lake has a long history of severe blue-green algae blooms. During the early 1980's, WDNR Bureau of Research conducted an experiment to determine if adding nitrogen to the lake would trigger a shift from nitrogen fixing blue-green algae species to non-bloom species (Lathrop 1988). The findings indicated that nitrogen applications were not effective due to short-term responses and other complicating factors. Since then, blue-green algae blooms in the lake have declined as a response to sustained dense aquatic plant growths and other factors.

Consistent with the hydrology of Fish and Crystal lakes, Indian Lake water levels have increased over time. The maximum recorded depth during the 1970s was 6 feet (Day et al. 1985). In 2006, the maximum water depth increased to 8.5 feet. The water levels in all three lakes may reflect increased regional groundwater recharge associated with agricultural conservation land use practices (Gebert and Krug 1996).

More recently, EWM and coontail were established in the lake and apparently suppressed phytoplankton blooms. Harvesting the dense beds became the primary management focus in the shallow lake. Dane County has been operating mechanical harvesters to create navigation channels for non-motorized boating access in the lake. These efforts also have potential to improve predator prey interactions.

Fish native species richness is limited by periodic past winterkills in the past. Species identified in the past surveys include fathead minnows (*Pimephales promelas*), bluntnose minnow (*Pimephales notatus*), white suckers (*Catostomus commersoni*), black bullhead (*Ameiurus melas*), green sunfish (*Lepomis cyanellus*), bluegill (*Lepomis macrochirus*), black crappie (*Pomoxis nigromaculatus*), largemouth bass (*Micropterus salmoides*), and yellow perch (*Perca flavescens*) (Lathrop 1988, Day et al. 1985). Following winterkills, bullhead populations periodically exploded and exacerbated turbidity and internal phosphorus loading in the lake. This occurred when dense bullhead populations disturbed bottom sediments while feeding. Currently, bluegill and largemouth bass populations are sustained by late winter aeration, and harvesting improves the habitat.

### **Recent Mechanical Harvesting Records**

Dane County records indicate that Indian Lake has been harvested since 1998. The primary goal is to create navigation channels within dense plant beds from the public boat ramp to various access points across the lake. Figure 11 displays annual tonnages of harvested aquatic plants removed from Indian Lake from 1989 to 2006, and Figure 12 shows Dane County harvesting priorities at Indian Lake.

Figure 11: Indian Lake mechanical harvesting summary





Figure 12: Indian Lake harvesting priorities for aquatic plants



## 2006 Aquatic Plant Survey Update

**Methods** (Refer to page 14)

### **Results and Discussion**

The aquatic plant survey was conducted on June 19, 2006. Secchi depth was measured at 4 feet with an equivalent Secchi TSI score of 54 (moderately eutrophic). The sampling grid contained 204 points (Figure 13), however the survey did not include a bay on the southeast side of the lake. Consistent with the Fish and Crystal lake surveys, the Indian Lake sampling grid did not reflect the higher water level and expanded lake area.

Very dense beds of coontail, CLP and EWM topped out of the water during the survey and rowing was difficult. CLP was beginning to decompose but still rigid enough to form dense beds. Coontail followed by EWM were the dominant two species yielding very high vegetated site and relative frequencies. Also abundant throughout the marshy lake were small floating plants, *Lemna and Wolffia*. Relative frequency data for both submersed and

floating plants are displayed in Figure 14. The average rake fullness was 2 for EWM and 3 for coontail. Beyond the sampling points, the entire lake was bordered by emergent vegetation, mostly cattails and to lesser extent bur reeds. Aquatic plant distributions are displayed in the Appendix G maps for the individual species. Appendix E notes fish and waterfowl values of desirable native plants in Fish, Crystal and Indian Lakes.

Figure 13: Point intercept sampling grid for Indian Lake





Figure 14: Indian Lake vegetated site occurrence and relative frequency results



The relative frequency describes the percentage that each aquatic plant species contributes to the whole aquatic plant community. Relative frequency is a commonly used metric since survey results can be compared with surveys that used different sampling techniques.

## Aquatic Plant Management Alternatives

Indian Lake offers outdoor enthusiasts a relatively rare opportunity to enjoy an undeveloped lake in southern Wisconsin. However, the dense aquatic plant communities have created nuisance conditions that impede navigation and detract somewhat from the experience. In the past, harvesting channels have improved boating access and benefited fish populations as well. Indian Lake has been suggested as a site for experimental whole-lake chemical treatment of EWM. This effort could tip the balance toward blue-green algae blooms. In 2006, EWM was not the dominant plant causing navigational problems. Rather, native coontail was more abundant. The existing conditions reflect decades of runoff pollution and sedimentation. Very high productivity, typical of hypereutrophic conditions, will likely persist in the lake. These conditions will continue unless a major rehabilitation project is launched, such as drawdown and dredging.

Mechanical harvesting often removes juvenile panfish in the conveyor with the vegetation. However, WDNR biologists consider the inadvertent removal of small panfish to have no effect on population levels. In some cases, removal of juvenile panfish may actually improve size structure when population levels are too high.

### Specific Options

- 1) **No treatment:** Rejecting all types of aquatic plant management is unreasonable given the high densities of EWM, CLP and coontail that undermine public access.
- 2) **Biological control:** This method does not appear realistic at this time. Weevil (*Euhrychiopsis lecontei*) populations had been well documented in the Fish Lake yet the density of EWM declined significantly only once over the past two decades. The findings are not clear if weevils were indeed responsible for the temporary decline in 1994. In Indian Lake, coontail and CLP currently combine for greater ecological and nuisance problems than EWM.
- 3) **Chemical control:** Herbicides are an unlikely management tool given the entire lake is encompassed within a park. An experimental whole-lake treatment may create severe blue-green algae blooms given the hypereutrophic status of the lake.

- 4) **Manual - hand removal:** Manually removing plants around piers and swimming areas is a viable option. Volunteers could be encouraged to remove dense weedy plants to create small fishing or swimming areas.
- 5) **Mechanical harvesting:** In 2006, mechanical harvesting is needed due to the high densities of coontail, EWM and CLP. Figure 12 demonstrates the areas typically harvested in Indian Lake most years.
- 6) **Physical control:** Most physical methods of controlling aquatic plants require permits from WDNR under Chapter 30 Wisconsin State Statutes. Hydraulic dredging is an unlikely option for Indian Lake. Sheets of dark fabrics anchored to the bottom covering the plants from any light create bottom or benthic barriers. A benthic barrier eliminates all plants, including non-target species, with the exception of free-floating plants and algae. These types of barriers can interfere with fish spawning and other pond wildlife. Over time, barriers may require maintenance due to siltation, ice damage, bubbling up and normal wear and tear. Fabrics applied in this way will require a permit. Fabrics are rarely used because of the labor of installation and maintenance. In local demonstrations, problems arose due to gas collection under the fabric. Drawdown combined with dredging could be evaluated but the option would be very costly and have dramatic effects on the ecosystem. Watershed sources of nutrients would also have to be controlled for an effective lake restoration effort.

## References

- Borman, S., R. Korth and J. Temte. 1997. Through the looking glass: a field guide to aquatic plants. WDNr Pub FH-207-97.
- Creed, R.P. 1998. A biogeographic perspective on Eurasian watermilfoil declines: additional evidence for the role of herbivorous weevils in promoting declines? *J. Aquat. Plant Manage.* 36:16-22.
- Cooke, D.G., E.B. Welch, S.A. Peterson and P.R. Newroth. 2005. Restoration and management of lakes and reservoirs. Third edition. Taylor and Francis/CRC Press. 616 p.
- DCRPC. 1979. Dane County Water Quality Plan.
- Day, E.A., G.P. Grzebieneak, K.M. Osterby, C.L. Brynildson. 1985. Surface water resources of Dane County. WDNr lake and stream classification project. Second edition. p. 15-17.
- Engel, S. 1987. The restructuring of littoral zones. *Lake Reserv. Manage.* 3: 235-242.
- Fassett, N.C. 1975. A manual of aquatic plants. The University of Wisconsin Press.
- Gebert, W.A. and W.R. Krug. 1996. Streamflow trends in Wisconsin's Driftless Area. *J. American Water Resources Association.* 32: 733-744.
- Janecek, J.A. 1988. Literature review on fishes interactions with aquatic macrophytes with special reference to the Upper Mississippi River System. U.S. Fish and Wildlife Service. Upper Mississippi River Conservation Committee. Rock Island, Illinois.
- Krohelski, J.T., Y. Lin, W.J. Rose and R.J. Hunt. 2002. Simulation of Fish, Mud, and Crystal Lakes and the shallow ground-water system, Dane County, Wisconsin. U.S. Geological Survey. Water Resources Investigations Report 02-4014.
- Larson, T. - personal communication, 10-06.
- Lathrop, R.C. 1988. Evaluation of whole-lake nitrogen fertilization for controlling blue-green algal blooms in a hypereutrophic lake. *Can. J. Fish. Aquat. Sci.* 45:2061-2075.
- Lillie, R.A. 2000. Temporal and spatial changes in milfoil distribution and biomass with weevils in Fish Lake, WI. *J. Aquat. Plant Manage.* 38:2000.
- Lillie, R.A. 1996. A quantitative survey of the floating-leafed and submersed macrophytes of Fish Lake, Dane County, Wisconsin. *Transactions Wisconsin Academy of Sciences, Arts and Letters.* 84:111-125.
- Lyons, J. 1992. Using the index of biotic integrity (IBI) to measure environmental quality in warmwater streams of Wisconsin. USDA General Technical Report NC-149.
- Madsen, J.D., J.W. Sutherland, J.A. Bloomfield, L.W., L.W. Eichler and C.W. Boylen. 1991. The decline of native vegetation under dense Eurasian watermilfoil canopies. *J. Aquat. Plant Manage.* 29: 94-99.
- Marshall, D.W., N.P. Nibbelink, P.J. Garrison, J. Panuska and S.R. Stewart. 1996. A management plan to protect and improve the Fish Lake ecosystem. U.S. EPA Clean Lakes Phase 1 Diagnostic and Feasibility Study report. 78 p.
- Marshall, D.W. 1990. Managing deep water stands of Eurasian watermilfoil to improve fish habitat and boating access. WDNr unpublished report.
- Mazzei, K.C., R.M. Newman, A. Loos and D. W. Ragsdale. 1999. Development rates of the native milfoil weevil, *Euhrychiopsis lecontei*, and damage to Eurasian watermilfoil at constant temperatures. *Biological Control* 16:139-143.
- Nichols, S.A. and J.G. Vennie. 1991. Attributes of Wisconsin Lake Plants. Wisconsin Geological and Natural History Survey. Information Circular 73.
- Olson, M.H., S.R. Carpenter, P. Cunningham, S. Gafny, B.R. Herwig, N.P. Nibbelink, T. Pellett, C. Storlie, A.S. Trebitz and K.A. Wilson. 1998. Managing macrophytes to improve fish growth: a multi-lake experiment. *Fisheries.* 23:6-12.

- Savino, J.F. and R.A. Stein. 1982. Predator-prey interaction between largemouth bass and bluegills as influenced by simulated, submersed vegetation. *Transactions of the American Fisheries Society*. 111: 255-265.
- Sheldon, S.P. and R.P. Creed. 1995. Use of native insect as a biological control for an introduced weed. *Ecological Applications* 5:1122-1132.
- Smith, C.S. and J.W. Barko. 1990. Ecology of Eurasian watermilfoil. *J. Aquat. Plant Manage.* 28:55-64.
- Trebitz, A., S. Carpenter, P. Cunningham, B. Johnson, R. Lillie, D. Marshall, T. Martin, R. Narf, T. Pellett, S. Stewart, C. Storlie and J. Unmuth. 1997. A model of bluegill-largemouth bass interactions in relation to aquatic vegetation and its management. *Ecological Modelling*. 94: 139-156.
- Unmuth, J.M., R.A. Lillie, D.S. Dreiksosen and D.W. Marshall. 2000. Influence of dense growth of Eurasian watermilfoil on lake water temperature and dissolved oxygen. *J. Freshwater Ecology*. 15:498-503.
- Unmuth, J.M. and M.J. Hansen. 1999. Effects of mechanical harvesting of Eurasian watermilfoil on largemouth bass and bluegill populations in Fish Lake, Wisconsin. *North American Journal of Fisheries Management*. 19:1089-1098.
- Wetzel, R.G. 1983. *Limnology*. Second Edition. Saunders College Publishing.

## GLOSSARY

<b>Alleopathy</b>	Harmful effect of one plant on another caused by the release of chemical compounds produced by the first plant.
<b>Biomanipulation</b>	A technique involving using predatory fish to reduce the number of fish that feed on organisms that eat algae.
<b>Chlorophyll a</b>	The green pigment present in all plant life and needed for photosynthesis. The amount present in lake water is related to the amount of algae found there and is used as an indicator of water quality.
<b>Columnaris</b>	Bacterial infection of fish which especially occurs when they are stressed. The disease is highly contagious to fish and typically enters through gills, mouth or small skin wounds.
<b>Cyanobacteria</b>	Another name for blue-green algae. A group of algae that are often associated with “problem” lake blooms. Certain species can produce toxins which can cause illness and even death in animals and humans. Blue-green algae can “fix” nitrogen from the atmosphere and thus are often found when phosphorus levels in water are high.
<b>Dessicated</b>	Something that is thoroughly dried.
<b>Emergent plants</b>	Species with leaves that extend above the water surface that are usually found in shallow water.
<b>Eutrophication</b>	The process by which lakes are enriched with nutrients thereby increasing the amount of rooted plants and algae. The extent to which this process has occurred is reflected in a trophic state classification system.
<b>Eutrophic</b>	Within a lake trophic state classification system, this is a lake that is rated as being very productive and fertile.
<b>Extirpation</b>	The act of being eliminated.
<b>Floating-leaf plants</b>	Rooted plants with leaves that float on the water surface, such as water lilies.
<b>Filamentous algae</b>	Algae that forms filaments or mats which attach to the bottom sediments, rooted plants, piers, etc.
<b>Hectare</b>	A metric unit of measure which is equivalent to about 2.47 acres.
<b>Herptiles</b>	A broad group of cold-blooded animals including turtles and amphibians.

<b>Hypereutrophic</b>	A very nutrient-enriched lake characterized by severe and dominant algal blooms and very poor water quality.
<b>Hypolimnion</b>	The layer of the lake closest to the bottom and immediately below the metalimnion or thermocline, within lakes deeper than 20 feet stratify, or layer, based on temperature differences.
<b>Hypolimnetic</b>	Referring to the hypolimnion.
<b>Internal loading</b>	Internal loading refers to sources of phosphorus within a lake, typically from deep water sediment or decaying aquatic plants in the littoral zone.
<b>Intolerant</b>	Aquatic species that are impacted by changing conditions. For example, if water quality worsens, certain intolerant species may disappear because of lowered oxygen levels or a loss of their habitat.
<b>Limnologist</b>	A specialist in the study of freshwater ponds and lakes.
<b>Littoral zone</b>	The shallow part of a lake where most of the rooted aquatic plants are found.
<b>Littoriprofundal</b>	Transition zone below the littoral zone, characterized by its lack of macrophytes; often adjacent to the metalimnion of stratified lakes.
<b>Macrophytes</b>	The higher (multi-celled) plants found growing in or near the water. They produce oxygen and provide food and cover for lake organisms.
<b>Mesotrophic</b>	Lakes that are in-between eutrophic (very fertile) and oligotrophic (infertile) waters. They exhibit fairly good water quality and rooted aquatic plant growth.
<b>Monotypic</b>	Involving only one species.
<b>Moss</b>	Mosses are bryophytes or non-vascular plants. These primitive plants lack flowers and seeds and produce spore capsules to reproduce. Mosses live in a variety of shady moist environments including deeper areas in clear lakes.
<b>Oligotrophic</b>	Very low nutrient, clear lakes having lower amounts of rooted aquatic plant growth and productivity, but rich in oxygen throughout their depth.
<b>Pelagic</b>	The open water zone of a lake outside of the littoral zone.

<b>Phosphorus entrainment</b>	Movement of phosphorus and other nutrients from the hypolimnion into the epilimnion due to wind-generated mixing and erosion of the thermocline.
<b>Phytoplankton</b>	Algae-like organisms found in waters that use light to support photosynthesis. Examples include diatoms, cyanobacteria and dinoflagellates.
<b>Planktivores</b>	A free-floating or drifting organism (example Daphnia) that feeds on phytoplankton.
<b>Planktonic algae</b>	Small free-floating algae including green algae, blue-green algae and diatoms.
<b>Point source</b>	Source of pollution (e.g. wastewater treatment plant) with a defined discharge point such as a discharge pipe.
<b>Secchi disc</b>	An 8-inch diameter disc with four alternating quadrants of black and white. It is lowered into a lake on a rope and used to measure light penetration. Lakes are infertile (oligotrophic) if the depth you can see the disc are great. Lakes are fertile (eutrophic) if the disc disappears quickly.
<b>Species richness</b>	The number of different species present.
<b>Thermocline</b>	The narrow transition zone between the epilimnion (top lake temperature layer) and the hypolimnion (bottom lake temperature layer). This portion of a lake is where the temperature changes most rapidly, and in most waters is found around 20 feet or deeper. Also called the metalimnion.
<b>Trophic State Index (TSI)</b>	A way to measure, rate and classify the quality of a water body. Trophic state (e.g. eutrophic, mesotrophic, oligotrophic and hypereutrophic) is a measure of biological productivity.
<b>Turion</b>	The over-wintering bud produced by aquatic plants.
<b>Two story fisheries</b>	Fisheries that support both warm water and cold water fish species, and that are thermally segregated for most of the year.
<b>Winter dormancy</b>	Refers to the condition of aquatic plants during the winter months, often in the form of seeds, turions, rhizomes or non-growing vegetation.